



Before



During



After



Summary

# *I dare* to learn all I can about starting treatment

## HERE'S WHAT TO EXPECT WHEN YOU START DARZALEX FASPRO<sup>®</sup>



### BEFORE

How to dress, and what to expect when you arrive at your healthcare provider's office.



### DURING

A quick description of the injection process for DARZALEX FASPRO<sup>®</sup>, and how long it takes.



### AFTER

What your healthcare provider may look for after your first few injections, and the importance of continuing your treatment as directed by your doctor.

## What is DARZALEX FASPRO<sup>®</sup> (daratumumab and hyaluronidase-fihj)?

DARZALEX FASPRO<sup>®</sup> is a prescription medicine used to treat adult patients with multiple myeloma:

- in combination with the medicines bortezomib, lenalidomide, and dexamethasone in newly diagnosed people who are eligible to receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib, lenalidomide, and dexamethasone in newly diagnosed people who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib, melphalan, and prednisone in people with newly diagnosed multiple myeloma who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines lenalidomide and dexamethasone in people with newly diagnosed multiple myeloma who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant) and in people whose multiple myeloma has come back or did not respond to treatment who have received at least one prior medicine to treat multiple myeloma
- in combination with the medicines bortezomib, thalidomide, and dexamethasone in newly diagnosed people who are eligible to receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib and dexamethasone in people who have received at least one prior medicine to treat multiple myeloma
- in combination with the medicines pomalidomide and dexamethasone in people who have received at least one prior medicine, including lenalidomide and a proteasome inhibitor, to treat multiple myeloma
- in combination with the medicines carfilzomib and dexamethasone in people whose multiple myeloma has come back or did not respond to treatment who have received one to three prior medicines to treat multiple myeloma
- alone in people who have received at least three prior medicines, including a proteasome inhibitor and an immunomodulatory agent, or did not respond to a proteasome inhibitor and an immunomodulatory agent

It is not known if DARZALEX FASPRO<sup>®</sup> is safe and effective in children.

## SELECT IMPORTANT SAFETY INFORMATION

**Do not receive DARZALEX FASPRO<sup>®</sup>** if you have a history of a severe allergic reaction to daratumumab, hyaluronidase, or any of the ingredients in DARZALEX FASPRO<sup>®</sup>. [Click here](#) for a complete list of ingredients in DARZALEX FASPRO<sup>®</sup>.

## BEFORE YOUR INJECTION

If this is the first time you're receiving an injection of DARZALEX FASPRO<sup>®</sup>, you may have some questions about what it's like and what you need to do to prepare. Here is some information that may help.

### PREPARING FOR THE INJECTION



#### Dress for comfort

Wear clothing that is loose around the waist.

DARZALEX FASPRO<sup>®</sup> is injected about 3 inches to the left or right of the belly button.



#### Set aside enough time

- To ask questions and discuss treatment goals
- To receive premedications and lab work if needed
- For your healthcare provider to monitor for a reaction to the injection, particularly for the first few injections that you receive

### BEFORE YOUR INJECTION

You will be given medicines to help reduce the risk of side effects to the injection, such as:



**Antihistamines** to prevent an allergic reaction



**Acetaminophen** or similar medicine to reduce fever



**Corticosteroids** to prevent inflammation



You may be given a quick physical exam before the injection. This includes **checking your pulse and blood pressure.**



**You will need to inform your healthcare providers and blood transfusion centers/personnel that you are taking DARZALEX FASPRO<sup>®</sup>.**

DARZALEX FASPRO<sup>®</sup> can affect the results of blood tests to match your blood type. These changes can last for up to 6 months after your final dose of DARZALEX FASPRO<sup>®</sup>. Your healthcare provider will help you with this and will do blood tests to match your blood type before you start treatment with DARZALEX FASPRO<sup>®</sup>. Tell all of your healthcare providers that you are being treated with DARZALEX FASPRO<sup>®</sup> before receiving blood transfusions if you need one.

### SELECT IMPORTANT SAFETY INFORMATION (cont)

**Before you receive DARZALEX FASPRO<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:**


- have a history of breathing problems
- have had shingles (herpes zoster)

Please [click here](#) to see the Product Information. Please [click here](#) to see additional Important Safety Information.




## DURING YOUR INJECTION

### PREPARATION



Your healthcare provider will prepare the syringe.

### LOCATION



An injection site on your stomach will be chosen and prepared. Injection sites in the stomach area will be rotated each time you receive treatment.

### TIMING



The injection will be completed in about 3 to 5 minutes.\*

\*This refers to the injection administration time and does not account for all aspects of treatment.

During your injection, your healthcare provider may use an injection needle or a winged infusion set.

injection  
needle



OR

winged  
infusion  
set



### SELECT IMPORTANT SAFETY INFORMATION (cont)

Before you receive DARZALEX FASPRO<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had or might now have a hepatitis B infection as DARZALEX FASPRO<sup>®</sup> could cause hepatitis B virus to become active again. Your healthcare provider will check you for signs of this infection before, during, and for some time after treatment with DARZALEX FASPRO<sup>®</sup>. Tell your healthcare provider right away if you get worsening tiredness or yellowing of your skin or white part of your eyes.

Please [click here](#) to see the Product Information. Please [click here](#) to see additional Important Safety Information.



## AFTER YOUR INJECTION





Pay attention to how you feel and tell your care team about any discomfort during or after treatment, and especially during the first and second injections. It could mean you may be having a reaction to the treatment.

**Do not receive DARZALEX FASPRO<sup>®</sup> if there is a history of a severe allergic reaction to daratumumab, hyaluronidase, or any of the ingredients in DARZALEX FASPRO<sup>®</sup>.**


Your healthcare provider may want you to remain in the office to watch for any side effects, serious allergic reactions, and other severe injection-related reactions. Serious allergic reactions and reactions due to release of certain substances by your body (systemic) that can lead to death can happen with DARZALEX FASPRO<sup>®</sup>.


Your healthcare provider may temporarily stop or completely stop treatment with DARZALEX FASPRO<sup>®</sup> if you have a serious reaction. Tell your healthcare provider or get medical help right away if you get any of these symptoms during or after an injection of DARZALEX FASPRO<sup>®</sup>.


 Shortness of breath or trouble breathing


 Dizziness or lightheadedness (hypotension)


 Cough

 Wheezing


 Heart beating faster than usual

 Low oxygen in the blood (hypoxia)

 Throat tightness or irritation

 Runny or stuffy nose

 Headache

 Itching

 High blood pressure

 Eye pain

 Nausea

 Vomiting


 Chills

 Fever


 Chest pain

 Blurred vision

### INJECTION SITE REACTIONS

 Skin reactions at or near the injection site (local), including injection site reactions, can happen with DARZALEX FASPRO<sup>®</sup>. Symptoms at the site of injection may include itching, swelling, bruising, pain, rash, bleeding, or redness of the skin. **These reactions sometimes happen more than 24 hours after an injection of DARZALEX FASPRO<sup>®</sup>.**

### POST-MEDICATION

 After your injection, you may also be given oral corticosteroids to reduce the risk of delayed reactions to DARZALEX FASPRO<sup>®</sup>.

### SELECT IMPORTANT SAFETY INFORMATION (cont)

**Before you receive DARZALEX FASPRO<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:**

- are pregnant or plan to become pregnant. DARZALEX FASPRO<sup>®</sup> may harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think that you may be pregnant during treatment with DARZALEX FASPRO<sup>®</sup>.
  - Females who are able to become pregnant should use an effective method of birth control (contraception) during treatment and for 3 months after your last dose of DARZALEX FASPRO<sup>®</sup>. Talk to your healthcare provider about birth control methods that you can use during this time.

Please [click here](#) to see the Product Information. Please [click here](#) to see additional Important Safety Information.



Before



During



After



Summary

## LOOKING FORWARD: HERE'S WHAT TO REMEMBER AS YOU CONTINUE ON DARZALEX FASPRO<sup>®</sup>



As you continue to receive DARZALEX FASPRO<sup>®</sup> injections in the long-term, your healthcare provider will develop a treatment plan that is right for you.

**Over time, DARZALEX FASPRO<sup>®</sup> treatments are needed only once every 4 weeks for most patients.**



**Make it a goal, and celebrate reaching a monthly dosing schedule.** By continuing treatment as directed by your doctor, you may give yourself the best chance of reaching the treatment goals discussed with your doctor.



### REMEMBER



**Dress comfortably** and in loose-fitting clothing.



**Be prepared for a quick physical exam** when you visit your healthcare provider.



You may need to take **medications before your injection** to help reduce the risk of side effects.



The injection time for DARZALEX FASPRO<sup>®</sup> is **about 3-5 minutes.\***

\*This refers to the injection administration time and does not account for all aspects of treatment.



Your healthcare provider may want you to remain in the office to **watch for any side effects, serious allergic reactions, and other severe injection-related reactions.**

DARZALEX withMe

### Personalized 1-on-1 Support

You have access to free, dedicated support. Your Care Navigator is here to help guide you to support solutions throughout your treatment journey.

Are you prescribed DARZALEX FASPRO<sup>®</sup>?

**Sign up for personalized support throughout your treatment journey now**

[Sign up for support](#) or call 833-565-9631, Monday through Friday, 8:00 AM-8:00 PM ET.

The support and resources provided by DARZALEX withMe are not intended to provide medical advice, replace a treatment plan you receive from your doctor or nurse, or serve as a reason for you to start or stay on treatment.

### SELECT IMPORTANT SAFETY INFORMATION (cont)

- Before starting DARZALEX FASPRO<sup>®</sup> in combination with lenalidomide, thalidomide, or pomalidomide, females and males must agree to the instructions in the lenalidomide, thalidomide, or pomalidomide REMS program.
- The lenalidomide, thalidomide, and pomalidomide REMS have more information about effective methods of birth control, pregnancy testing, and blood donation for females who can become pregnant.
- For males who have female partners who can become pregnant, there is information in the lenalidomide, thalidomide, and pomalidomide REMS about sperm donation and how lenalidomide, thalidomide, and pomalidomide can pass into human semen.

Please [click here](#) to see the Product Information. Please [click here](#) to see additional Important Safety Information.



Before



During



After



Summary

# INDICATIONS AND IMPORTANT SAFETY INFORMATION FOR DARZALEX FASPRO<sup>®</sup>

## What is DARZALEX FASPRO<sup>®</sup> (daratumumab and hyaluronidase-fihj)?

DARZALEX FASPRO<sup>®</sup> is a prescription medicine used to treat adult patients with multiple myeloma:

- in combination with the medicines bortezomib, lenalidomide, and dexamethasone in newly diagnosed people who are eligible to receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib, lenalidomide, and dexamethasone in newly diagnosed people who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib, melphalan, and prednisone in people with newly diagnosed multiple myeloma who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines lenalidomide and dexamethasone in people with newly diagnosed multiple myeloma who cannot receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant) and in people whose multiple myeloma has come back or did not respond to treatment who have received at least one prior medicine to treat multiple myeloma
- in combination with the medicines bortezomib, thalidomide, and dexamethasone in newly diagnosed people who are eligible to receive a type of stem cell transplant that uses their own stem cells (autologous stem cell transplant)
- in combination with the medicines bortezomib and dexamethasone

in people who have received at least one prior medicine to treat multiple myeloma

- in combination with the medicines pomalidomide and dexamethasone in people who have received at least one prior medicine, including lenalidomide and a proteasome inhibitor, to treat multiple myeloma
- in combination with the medicines carfilzomib and dexamethasone in people whose multiple myeloma has come back or did not respond to treatment who have received one to three prior medicines to treat multiple myeloma
- alone in people who have received at least three prior medicines, including a proteasome inhibitor and an immunomodulatory agent, **or** did not respond to a proteasome inhibitor and an immunomodulatory agent

It is not known if DARZALEX FASPRO<sup>®</sup> is safe and effective in children.

## IMPORTANT SAFETY INFORMATION

**Do not receive DARZALEX FASPRO<sup>®</sup>** if you have a history of a severe allergic reaction to daratumumab, hyaluronidase, or any of the ingredients in DARZALEX FASPRO<sup>®</sup>. See below for a complete list of ingredients in DARZALEX FASPRO<sup>®</sup>.

**Before you receive DARZALEX FASPRO<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:**

- have a history of breathing problems
- have had shingles (herpes zoster)
- have ever had or might now have a hepatitis B infection as

DARZALEX FASPRO<sup>®</sup> could cause hepatitis B virus to become active again. Your healthcare provider will check you for signs of this infection before, during, and for some time after treatment with DARZALEX FASPRO<sup>®</sup>. Tell your healthcare provider right away if you get worsening tiredness or yellowing of your skin or white part of your eyes.

- are pregnant or plan to become pregnant. DARZALEX FASPRO<sup>®</sup> may harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think that you may be pregnant during treatment with DARZALEX FASPRO<sup>®</sup>.
  - Females who are able to become pregnant should use an effective method of birth control (contraception) during treatment and for 3 months after your last dose of DARZALEX FASPRO<sup>®</sup>. Talk to your healthcare provider about birth control methods that you can use during this time.
  - Before starting DARZALEX FASPRO<sup>®</sup> in combination with lenalidomide, thalidomide, or pomalidomide, females and males must agree to the instructions in the lenalidomide, thalidomide, or pomalidomide REMS program.
    - The lenalidomide, thalidomide, and pomalidomide REMS have more information about effective methods of birth control, pregnancy testing, and blood donation for females who can become pregnant.
    - For males who have female partners who can become pregnant, there is information in the lenalidomide, thalidomide, and pomalidomide REMS about sperm donation and how lenalidomide, thalidomide, and pomalidomide can pass into human semen.



Before



During



After



Summary

## INDICATIONS AND IMPORTANT SAFETY INFORMATION FOR DARZALEX FASPRO<sup>®</sup> (cont)

- are breastfeeding or plan to breastfeed. It is not known if DARZALEX FASPRO<sup>®</sup> passes into your breast milk. You should not breastfeed during treatment with DARZALEX FASPRO<sup>®</sup>. Talk to your healthcare provider about the best way to feed your baby during treatment with DARZALEX FASPRO<sup>®</sup>.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

### How will I receive DARZALEX FASPRO<sup>®</sup>?

- DARZALEX FASPRO<sup>®</sup> may be given alone to treat multiple myeloma or together with other medicines used to treat multiple myeloma
- DARZALEX FASPRO<sup>®</sup> will be given to you by your healthcare provider as an injection under the skin in the stomach area (abdomen).
- DARZALEX FASPRO<sup>®</sup> is injected over 3 to 5 minutes.
- Your healthcare provider will decide the time between doses as well as how many treatments you will receive.
- Your healthcare provider will give you medicines before each dose of DARZALEX FASPRO<sup>®</sup> and after each dose of DARZALEX FASPRO<sup>®</sup> to help reduce the risk of serious allergic reactions and other reactions due to release of certain substances by your body (systemic).

If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment.

### DARZALEX FASPRO<sup>®</sup> may cause serious reactions, including:

- **Serious allergic reactions and other severe injection-related reactions.** Serious allergic reactions and reactions due to release of certain substances by your body (systemic) that can lead to death can happen with DARZALEX FASPRO<sup>®</sup>. Your healthcare provider may temporarily stop or completely stop treatment with DARZALEX FASPRO<sup>®</sup> if you have a serious reaction. Tell your healthcare provider or get medical help right away if you get any of these symptoms during or after an injection of DARZALEX FASPRO<sup>®</sup>.

- |  |                       |
|--|-----------------------|
| ◦ shortness of breath or trouble breathing   | ◦ headache            |
| ◦ dizziness or lightheadedness (hypotension) | ◦ itching             |
| ◦ cough                                      | ◦ high blood pressure |
| ◦ wheezing                                   | ◦ eye pain            |
| ◦ heart beating faster than usual            | ◦ nausea              |
| ◦ low oxygen in the blood (hypoxia)          | ◦ vomiting            |
| ◦ throat tightness or irritation             | ◦ chills              |
| ◦ runny or stuffy nose                       | ◦ fever               |
|  | ◦ chest pain          |
|  | ◦ blurred vision      |

- **Injection site reactions.** Skin reactions at or near the injection site (local), including injection site reactions, can happen with DARZALEX FASPRO<sup>®</sup>. Symptoms at the site of injection may include itching, swelling, bruising, pain, rash, bleeding, or redness of the skin. These reactions sometimes happen more than 24 hours after

an injection of DARZALEX FASPRO<sup>®</sup>.

- **Infections.** DARZALEX FASPRO<sup>®</sup> can cause serious life-threatening infections that can lead to death. Tell your healthcare provider right away if you develop a fever, trouble breathing, cough, burning or pain when you urinate, or any other signs and symptoms of infection during treatment.
- **Decreases in blood cell counts.** DARZALEX FASPRO<sup>®</sup> can decrease white blood cell counts, which help fight infections, and blood cells called platelets, which help to clot blood. Decreases in blood cell counts are common with DARZALEX FASPRO<sup>®</sup> and can be severe. Your healthcare provider will check your blood cell counts during treatment with DARZALEX FASPRO<sup>®</sup>. Tell your healthcare provider if you develop fever or have signs of bruising or bleeding.
- **Changes in blood tests.** DARZALEX FASPRO<sup>®</sup> can affect the results of blood tests to match your blood type. These changes can last for up to 6 months after your final dose of DARZALEX FASPRO<sup>®</sup>. Your healthcare provider will do blood tests to match your blood type before you start treatment with DARZALEX FASPRO<sup>®</sup>. **Tell all of your healthcare providers that you are being treated with DARZALEX FASPRO<sup>®</sup> before receiving blood transfusions.**



Before



During



After



Summary

## IMPORTANT SAFETY INFORMATION FOR DARZALEX FASPRO<sup>®</sup> (cont)

### The most common side effects of DARZALEX FASPRO<sup>®</sup> when used alone include:

- cold-like symptoms (upper respiratory tract infection)
- muscle, bone, and joint pain
- tiredness
- diarrhea
- rash
- trouble sleeping, including sleep apnea and restless legs
- nerve problems, including increased or decreased sensitivity to touch, temperature, or pain; loss of smell; and numbness, tingling, or burning sensations
- injection site reactions

### The most common side effects of DARZALEX FASPRO<sup>®</sup> when used in combination therapy include:

- tiredness
- nausea
- diarrhea
- shortness of breath
- trouble sleeping
- headache
- rash
- kidney problems
- movement and balance problems, muscle spasms, weakness, and tremors
- fever
- cough
- muscle spasms
- back pain
- vomiting
- high blood pressure
- muscle, bone, and joint pain
- decreased appetite
- urinary tract infection
- stomach pain
- cold-like symptoms (upper respiratory tract infection)
- nerve problems, including increased or decreased sensitivity to touch, temperature, or pain; loss of smell; and numbness, tingling, or burning sensations
- constipation
- lung infection (pneumonia)
- swollen hands, ankles, or feet
- feeling dizzy
- bruising
- COVID-19

The most common severe abnormal blood test results with DARZALEX FASPRO<sup>®</sup> include decreased white blood cells, platelets, and red blood cells.

These are not all of the possible side effects of DARZALEX FASPRO<sup>®</sup>. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### General information about the safe and effective use of DARZALEX FASPRO<sup>®</sup>

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your pharmacist or healthcare provider for information about DARZALEX FASPRO<sup>®</sup> that is written for health professionals.

**Active ingredient:** daratumumab and hyaluronidase-fihj

**Inactive ingredients:** L-histidine, L-histidine hydrochloride monohydrate, L-methionine, polysorbate 20, sorbitol, water for injection

Please [click here](#) to read the full Prescribing Information for DARZALEX FASPRO<sup>®</sup>.

cp-143282v13